arXiv:1703.05817v1 [astro-ph.SR] 16 Mar 2017

NEAR-IR TRIGONOMETRIC PARALLAXES OF NEARBY STARS IN THE GALACTIC PLANE USING THE VVV [9]

J. C. Beamín[1,2], R. A. Mendez[3,2], R. L. Smart[4], R. Jara[3], R. Kurtev[1,2], M. Gromadzki[5], V. Villanueva[1], D. Minniti[6,2,7], L.C. Smith[8], P. W. Lucas[8]

RESUMEN

ABSTRACT

We use the multi-epoch K_S band observations, covering a ∼5 years baseline to obtain milli and sub-milli arcsec precision astrometry for a sample of eighteen previously known high proper motion sources, including precise parallaxes for these sources for the first time. In this study we show the capability of the VVV project to measure high precision trigonometric parallaxes for very low mass stars (VLMS) up to distances of ∼250 pc reaching farther than most other ground based surveys or space missions for these types of stars. Additionally, we used spectral energy distribution to search for evidence of unresolved binary systems and cool sub-dwarfs. We detected five systems that are most likely VLMS belonging to the Galactic halo based on their tangential velocities, and four objects within 60 pc that are likely members of the thick disk. A more comprehensive study of high proper motion sources and parallaxes of VLMS and brown dwarfs with the VVV is ongoing , including thousands of newly discovered objects.

Key Words: Methods: Astrometry — Stars: low mass — Stars: Brown dwarfs — Stars: subdwarfs — Methods: Proper motions

1. INTRODUCTION

The VVV survey is a multi-epoch near infrared ESO public survey using the VISTA 4.2 m telescope, observing over 560 sq. degrees towards the most crowded regions of the sky, the galactic bulge and disk. VVV mapped these areas in five NIR bands (ZY JHKs) and had a variability campaign lasting 5-6 years in the Ks band, obtaining between ∼70 and ∼150 epochs per pointing. Additionally, the small pixel size of 0.34" (and a typical FWHM of 1" in the Ks band) provide a good resolution and make the VVV survey ideal to find new moving sources in the most crowded areas in the sky.

Taking advantage of all these characteristics we performed a search for high proper motions sources and we found thousands of them (Kurtev et al. 2017; Smith, L.C. 2016).

We were able to obtain spectroscopic follow up observations for several tens of sources and the results were discussed in Beamín et al. 2015 and Gromadzki et al. 2016, most of them K-M dwarfs. Additionally a new sample of ∼50 ultra cool dwarfs (UCDs) with full atrometric characterization and classification with low-mid Near IR resoluion spectra from the VVV survey is being analyzed and made publicly available soon (Beamin et. al. in prep).

We selected sources from previous studies of high proper motion (HPM), lying within the VVV area that were not saturated on the images. We obtained ten sources from the studies of Lépine (2005) and Lépine (2008), and seven co-moving companions discovered by Ivanov et al. (2013), and VVV BD001 Beamín et al. (2013)

Among our discoveries two of them are particularly interesting an unusually blue L dwarf at 18.6 pc (revised distance, this work) and a benchmark early L dwarf companion to the ∼400 Myr star β Circini (Smith et al. 2015).

In this work we focus in obtaining precise parallax measurements for a sample of cool stars within 250 pc from the sun. Some of the stars we selected

[1]Instituto de Física y Astronomía, Facultad de Ciencias, Universidad de Valparaíso, Ave. Gran Bretaña 1111, Playa Ancha, Valparaíso, Chile
[2]Millennium Institute of Astrophysics, Chile.
[3]Universidad de Chile, Departamento de Astronomía, Casilla 36-D, Santiago, Chile.
[4]INAF-Osservatorio Astrofisico di Torino, Strada Osservatorio 20, 10025 Pino Torinese, TO, Italy.
[5]Warsaw University Astronomical Observatory, Al. Ujazdowskie 4, 00-478 Warszawa, Poland
[6]Facultad de Ciencias Exactas, Universidad Andres Bello, Fernandez Concha 700, Las Condes, Santiago, Chile
[7]Vatican Observatory, V00120 Vatican City State, Italy
[8]Centre for Astrophysics Research, Science and Technology Research Institute, University of Hertfordshire, Hatfield AL10 9AB, UK
[9]BASED ON OBSERVATIONS TAKEN WITHIN THE ESO VISTA PUBLIC SURVEY VVV, PROGRAMME ID 179.B-2002

will also be observed by Gaia, allowing us to cross-check the astrometric accuracy of the VVV survey.

2. METHDS

2.1. *Target Selection*

Target selection: We selected sources from previous studies of high proper motion (HPM), particularly, we selected all the sources in the VVV area that were not saturated in the VVV images. We obtained ten sources from the studies of Lépine 2005 and Lépine 2008, seven co-moving companions discovered by Ivanov et al. 2013, and VVV BD001 from Beamín et al. 2013.

2.2. *Astrometric procedure*

For these eighteen sources we selected reference stars in a circle of radius 1' around the target which, given the high stellar density in these fields, provides an adequate number of reference stars (above a hundred sources). These were selected among the highest S/N objects in the field of view, satisfying the condition that they appear in at least 80% of the frames, and do not exhibit large proper motions.

After all epochs were on the same astrometric reference system, we fit proper motion, parallax and a shift per coordinate. More details on the procedure can be found in Smart et al. (2003) and Beamín et al. (2015).

3. RESULTS

We obtained improved proper motion and parallaxes for the first time for these 17 sources, also provide revised proper motion and parallax for an additional source (VVV BD 001). In Fig. 1 we show a comparison between our estimates and previous results of proper motion in RA and DEC. The green points show stars from Ivanov et al. 2013 and blue dots stars from Lépine 2005, 2008, and the red dot is VVVBD-001. The red solid line show a 1:1 relation, the black dashed line is the best fit with blue shade marking 1-σ interval. We reach milli and sub-milli arcsec precision in a couple of sources,comparable with space based observations and results from state of the art ground based astrometric projects. A summary of our astrometric results is shown in Table 3.

3.1. *Remarkable objects*

2MASS J13124062-6408313 is a wide binary most likely a member of the Halo, given the colours, and large tangential velocity (Vtan >300 Km/s), there are another five sources that have tangential velocities greater than 200 km/s making them most likely



Fig. 1. Comparison of our measurements of proper motion in right ascension (upper panel) and declination (lower panel) with the values listed in the literature. Green dots are from Ivanov et al. 2013, blue dots from Lépine 2005, 2008, and the red dot is VVV BD 001 (Beamín et al. 2013). The error bars of each object are smaller than the point size

members of the galactic halo, most of them also show unusual colours making them good candidates for subdwarf or unresolved binaries We tested our parallax against the new Tycho-Gaia astrometric solution (TGAS) (Gaia Collaboration et al. 2016), in the catalog we found parallaxes for L 149-77 and L 200-41 and we present parallaxes for L 149-77 B, L 200-41 B. The result are in agreement within 1.5σ, 1σ with the Gaia results for the primaries. Examples of the source motion, including parallax for the nearest and farthest away objects in our study are shown in Fig 2. Finally, 2MASS J17433729-2303586 has a small chance to produce a microlensing event in the next 5 years, a detailed analysis and predictions for this source are being carried on.

4. CONCLUSIONS

We present improved NIR astrometry including parallaxes for eighteen sources. We found five sources that are more likely nearby halo objects, in

Name	π	σ_π	$\mu_\alpha \cos\delta$	$\sigma_{\mu_\alpha cos\,\delta}$	μ_δ	σ_{μ_δ}	K_S
	mas	mas	mas yr^{-1}	mas yr^{-1}	mas yr^{-1}	mas yr^{-1}	mag
2MASS J13124062-6408313 B	7.66	2.88	-494.80	1.90	61.10	1.41	13.297
2MASS J13314733-6305061	7.08	3.25	-571.92	1.81	108.30	1.92	12.917
L 149-77 B	22.20	2.31	173.25	1.30	193.46	1.11	10.418
LHS 2881 B	37.87	2.08	-537.41	1.10	-537.68	1.20	10.863
2MASS J14570297-5943468	15.44	1.67	-388.77	0.87	-472.73	0.96	11.780
L 200-41 B	8.53	2.57	-169.60	1.52	-142.38	1.73	10.280
2MASS J15291530-5620406	12.58	2.89	-583.73	2.00	-129.01	1.81	12.480
2MASS J15495475-5524582	17.69	1.64	-521.57	0.82	-273.14	0.97	11.753
VVV J172640.22-273803.13	53.93	1.20	-546.38	0.74	-322.29	0.82	12.194
2MASS J17433729-2303586	9.59	0.76	-86.19	0.60	-477.57	0.75	11.031
2MASS J17442929-2044481	25.91	2.10	-242.02	1.60	-689.99	1.90	10.599
2MASS J17505469-2734335	14.55	1.60	-397.35	1.00	-358.95	1.00	10.947
2MASS J17512332-3505258	3.91	0.98	-150.74	0.60	-129.30	0.60	11.339
2MASS J17521105-3916546	18.04	1.22	-88.90	0.82	-367.84	0.89	10.988
MACHO 124.22158.2900	9.55	1.22	-143.11	0.91	-245.84	0.89	11.562
MACHO 124.22158.2910	5.91	1.71	-141.50	1.30	-246.39	1.11	12.09
J18092457-4104351	31.01	2.65	173.58	1.44	-386.85	1.83	10.608
		$<\sigma_\pi>$		$<\sigma\mu_\alpha>$		$<\sigma\mu_\delta>$	
		1.92		1.2		1.23	

TABLE 1

ASTROMETRY FOR THE EIGHTEEN SOURCES IN THIS STUDY LAST ROW SHOWS THE AVERAGE OF THE ERRORS IN PARALLAX AND PROPER MOTION AT EACH COORDINATE.

particular two wide binary system, and one probable unresolved binary. Spectroscopic follow up for these sources is encouraged, particularly RV measurements for the wide binaries with high tangential velocity. Our results show consistent distances for sources up to distances of ∼250 pc. And in the case of co-moving companions our parallax measurements agree within the errors with the parallax calculated for the primary source by the Gaia mission. Finally we found one source that might produce a microlensing event in the upcoming years, we encourage high spatial resolution imaging, spectroscopic and time series photometric follow up for this target. *Acknowledgments This study was supported by programa Comité Mixto ESO-Gobierno de Chile. This publication makes use of data products from the Two Micron All Sky Survey, which is a joint project of the University of Massachusetts and the Infrared Processing and Analysis Center/California Institute of Technology, funded by the National Aeronautics and Space Administration and the National Science Foundation, and SIMBAD database, operated at CDS, Strasbourg, France.*

REFERENCES

Beamín, J. C., Minniti, D., Gromadzki, M., et al. 2013, A&A*, 557, L8*

Beamín, J. C., Ivanov, V. D., Minniti, D., et al. 2015, MNRAS*, 454, 4054*

Gaia Collaboration, Prusti, T., de Bruijne, J. H. J., et al. 2016, A&A*, 595, A1*

Gromadzki, M., Kurtev, R., Beamin, J. C., et al. 2016, AcA, 66, 293

Ivanov, V. D., Minniti, D., Hempel, M., et al. 2013, A&A*, 560, A21*

Kurtev, R., Gromadzki, M., Beamín, J. C., et al. 2017, MNRAS*, 464, 1247*

Lépine, S. 2005, AJ*, 130, 1247*

Lépine, S. 2008, AJ*, 135, 2177*

Minniti, D., Lucas, P. W., Emerson, J. P., et al. 2010, NewA, 15, 433

Smart, R. L., Lattanzi, M. G., Bucciarelli, B., et al. 2003, A&A*, 404, 317*

Smith, L. C., Lucas, P. W., Contreras Peña, C., et al. 2015, MNRAS*, 454, 4476*

Smith, L. C., 2016, PhD. Thesis


Relative Y Postion [arcseconds]
Relative X Position [arcseconds]
Relative Y Postion [arcseconds]
Relative X Position [arcseconds]

Fig. 2. Motion on the sky for the closest and farthest source in this study and the best fit, including proper motion and parallax. Upper panel, VVV BD001, the estimated distance is 18.55 $\pm^{0.42}_{0.4}$ pc. Lower panel 2MASS J17512332-3505258, located at a distance of 255$\pm^{86}_{51}$ pc.